



## Advances in the Theory of Nonlinear Analysis and its Applications

ISSN: 2587-2648

Peer-Reviewed Scientific Journal

# Existence of positivity of the solutions for higher order three-point boundary value problems involving $p$ -Laplacian

Sreedhar Namburi<sup>a</sup>, Ravi Sankar Ronanki<sup>a</sup>, Rajendra Prasad Kapula<sup>b</sup>

<sup>a</sup>Department of Mathematics, GITAM (Deemed to be University), Visakhapatnam, India.

<sup>b</sup>Department of Applied Mathematics, Andhra University, Visakhapatnam, India.

## Abstract

The present study focusses on the existence of positivity of the solutions to the higher order three-point boundary value problems involving  $p$ -Laplacian

$$\begin{aligned} [\phi_p(x^{(m)}(t))]^{(n)} &= g(t, x(t)), \quad t \in [0, 1], \\ x^{(i)}(0) &= 0, \quad \text{for } 0 \leq i \leq m-2, \\ x^{(m-2)}(1) - \alpha x^{(m-2)}(\xi) &= 0, \\ [\phi_p(x^{(m)}(t))]_{\text{at } t=0}^{(j)} &= 0, \quad \text{for } 0 \leq j \leq n-2, \\ [\phi_p(x^{(m)}(t))]_{\text{at } t=1}^{(n-2)} - \alpha [\phi_p(x^{(m)}(t))]_{\text{at } t=\xi}^{(n-2)} &= 0, \end{aligned}$$

where  $m, n \geq 3$ ,  $\xi \in (0, 1)$ ,  $\alpha \in (0, \frac{1}{\xi})$  is a parameter. The approach used by the application of Guo-Krasnosel'skii fixed point theorem to determine the existence of positivity of the solutions to the problem.

**Keywords:** three-point, nonlinear, boundary value problem,  $p$ -Laplacian, Green's function, positive solution.

**2010 MSC:** 34B15, 34A40.

---

Email addresses: [sreedharnamburi13@gmail.com](mailto:sreedharnamburi13@gmail.com) (Sreedhar Namburi), [ronanki1980@gmail.com](mailto:ronanki1980@gmail.com) (Ravi Sankar Ronanki), [rajendra92@rediffmail.com](mailto:rajendra92@rediffmail.com) (Rajendra Prasad Kapula)

Received December 22, 2020, Accepted June 22, 2022, Online June 24, 2022

## 1. Introduction

Differential equations have created a tremendous amount of interest and played a vital role in many areas of mathematical sciences. The theory of differential equations gives profound and wide mathematical support for addressing many emerging issues of present society that are challenging and multidisciplinary in the universe. In this theory, one of the significant and useful operators is one-dimensional  $p$ -Laplacian operator and is defined as  $\phi_p(\tau) = |\tau|^{p-2}\tau$ , where  $p > 1$ ,  $\phi_p^{-1} = \phi_q$  and  $\frac{1}{p} + \frac{1}{q} = 1$ . Recently, researchers have given great consideration to study  $p$ -Laplacian problems due to the wide applicability in various real time applications such as biophysics, plasma physics, image processing, rheology, glaciology, turbulent filtration in porous media, radiation of heat etc. We mention a few papers devoted to  $p$ -Laplacian problems, see [3, 4, 19, 26, 20, 10, 13] for the existence of positivity of the solutions. For applications and recent developments, we refer [7, 1, 2, 5, 9, 15, 16, 18, 28].

We consider higher order three-point boundary value problems involving  $p$ -Laplacian of the form

$$[\phi_p(x^{(m)}(t))]^{(n)} = g(t, x(t)), \quad t \in [0, 1], \quad (1)$$

$$\left. \begin{aligned} x^{(i)}(0) &= 0, \text{ for } 0 \leq i \leq m-2, \\ x^{(m-2)}(1) - \alpha x^{(m-2)}(\xi) &= 0, \\ [\phi_p(x^{(m)}(t))]_{\text{at } t=0}^{(j)} &= 0, \text{ for } 0 \leq j \leq n-2, \\ [\phi_p(x^{(m)}(t))]_{\text{at } t=1}^{(n-2)} - \alpha [\phi_p(x^{(m)}(t))]_{\text{at } t=\xi}^{(n-2)} &= 0, \end{aligned} \right\} \quad (2)$$

where  $m, n \geq 3$ ,  $\xi \in (0, 1)$ ,  $\alpha \in (0, \frac{1}{\xi})$  is a parameter, and the function  $g : [0, 1] \times \mathbb{R}^+ \rightarrow \mathbb{R}^+$  is continuous and establish the existence of positivity of the solution by using fixed point theorem of Guo–Krasnosel'skii. If  $p = 2$ , we get various order three-point boundary value problems by giving different values to  $m$  and  $n$ . In the past, most researchers have focussed and demonstrated the positivity results for boundary value problems of third order three-point using various methods, see [11, 29, 33, 35, 22, 23, 25, 27, 34, 24, 39]. However, some works on positivity results have been found for  $n^{\text{th}}$ ,  $2n^{\text{th}}$  and  $3n^{\text{th}}$  order  $p$ -Laplacian boundary value problems, see [21, 12, 32, 38, 8, 36, 37, 30, 31]. Motivated by the aforementioned papers, we then extend the results to  $mn^{\text{th}}$  order  $p$ -Laplacian problem stated in (1), (2).

For establishing the results, assume the following condition is fulfilled in the entire paper:

(F1)  $\alpha$  is a parameter such that  $0 < \alpha\xi < 1$ , where  $\xi \in (0, 1)$ .

The remaining portion of the paper is structured as below. With the aid of Green functions, the solution of  $p$ -Laplacian problem stated in (1) and (2) is expressed as a solution to an analogous integral equation and then some inequalities for these Green functions are established in Section 2. The existence of the positivity results of the problem (1)-(2) is established and the results are validated by an example in Section 3.

## 2. Preliminaries

This section contains preparatory results that are necessary to demonstrate the existence results.

We first express the solution of the following  $m^{\text{th}}$  order non-homogeneous problem of three-point

$$x^{(m)}(t) + \varphi(t) = 0, \quad t \in [0, 1], \quad (3)$$

$$\left. \begin{aligned} x^{(i)}(0) &= 0, \text{ for } 0 \leq i \leq m-2, \\ x^{(m-2)}(1) - \alpha x^{(m-2)}(\xi) &= 0, \end{aligned} \right\} \quad (4)$$

where  $\varphi(t) \in C([0, 1], \mathbb{R}^+)$ , in terms of Green's function  $\overline{G}_m(t, s)$  as a solution of an analogous integral equation. By taking  $y(t) = \phi_p(x^{(m)}(t))$ , the solution of  $n^{\text{th}}$  order non-homogeneous problem of three-point

$$y^{(n)}(t) + \psi(t) = 0, \quad t \in [0, 1], \quad (5)$$

$$\left. \begin{aligned} y^{(j)}(0) &= 0, \text{ for } 0 \leq j \leq n-2, \\ y^{(n-2)}(1) - \alpha y^{(n-2)}(\xi) &= 0, \end{aligned} \right\} \quad (6)$$

where  $\psi(t) \in C([0, 1], R^+)$ , is expressed in terms of Green's function  $G_n(t, s)$  as a solution of an analogous integral equation.

**Lemma 2.1.** [34] *If the condition (F1) is fulfilled, then the solution of the problem stated in (3), (4) is*

$$x(t) = \int_0^1 \overline{G}_m(t, s) \varphi(s) ds,$$

where

$$\overline{G}_m(t, s) = H(t, s) + \frac{\alpha t^{m-1}}{(m-1)! (1-\xi\alpha)} G(\xi, s), \quad (7)$$

$$H(t, s) = \frac{1}{(m-1)!} \begin{cases} t^{m-1}(1-s), & 0 \leq t \leq s \leq 1, \\ [t^{m-1}(1-s) - (t-s)^{m-1}], & 0 \leq s \leq t \leq 1, \end{cases}$$

and

$$G(\xi, s) = \begin{cases} (1-\xi)s, & 0 \leq s \leq \xi \leq 1, \\ (1-s)\xi, & 0 \leq \xi \leq s \leq 1. \end{cases}$$

**Lemma 2.2.** [34] *If the condition (F1) is fulfilled, then the solution of the problem stated in (5), (6) is*

$$y(t) = \int_0^1 G_n(t, s) \psi(s) ds,$$

where

$$G_n(t, s) = K(t, s) + \frac{\alpha t^{n-1}}{(n-1)! (1-\xi\alpha)} G(\xi, s), \quad (8)$$

$$K(t, s) = \frac{1}{(n-1)!} \begin{cases} t^{n-1}(1-s), & 0 \leq t \leq s \leq 1, \\ [t^{n-1}(1-s) - (t-s)^{n-1}], & 0 \leq s \leq t \leq 1, \end{cases}$$

and

$$G(\xi, s) = \begin{cases} (1-\xi)s, & 0 \leq s \leq \xi \leq 1, \\ (1-s)\xi, & 0 \leq \xi \leq s \leq 1. \end{cases}$$

Using Lemmas 2.1 and 2.2, the solution of the problem stated in (1), (2) is

$$x(t) = \int_0^1 \overline{G}_m(t, s) \phi_q \left[ \int_0^1 G_n(s, r) g(r, x(r)) dr \right] ds. \quad (9)$$

**Lemma 2.3.** [34] *If the condition (F1) is fulfilled, then  $\overline{G}_m(t, s)$  in (7) fulfills the subsequent conditions:*

- (i)  $\overline{G}_m(t, s) \geq 0$ , for all  $t \in [0, 1]$  and  $s \in [0, 1]$ ,
- (ii)  $\overline{G}_m(t, s) \leq \overline{G}_m(1, s)$ , for all  $t \in [0, 1]$  and  $s \in [0, 1]$ ,
- (iii)  $\min_{t \in [\xi, 1]} \overline{G}_m(t, s) \geq \xi^{m-1} \overline{G}_m(1, s)$ , for all  $s \in [0, 1]$ .

**Lemma 2.4.** [34] *If the condition (F1) is fulfilled, then  $G_n(t, s)$  in (8) fulfills the subsequent conditions:*

- (i)  $G_n(t, s) \geq 0$ , for all  $t \in [0, 1]$  and  $s \in [0, 1]$ ,

(ii)  $G_n(t, s) \leq G_n(1, s)$ , for all  $t \in [0, 1]$  and  $s \in [0, 1]$ ,

(iii)  $\min_{t \in [\xi, 1]} G_n(t, s) \geq \xi^{n-1} G_n(1, s)$ , for all  $s \in [0, 1]$ .

The fixed point theorem of Guo–Krasnosel'skii mentioned below is often used as the fundamental tool to establish the positivity results of the problem stated in (1), (2).

**Theorem 2.5.** [6, 14, 17] Let  $X$  be a Banach Space and the set  $\kappa \subseteq X$  be a cone. Suppose the sets  $\Omega_1$  and  $\Omega_2$  are any two open subsets of  $X$  such that  $0 \in \Omega_1$  and  $\overline{\Omega}_1 \subset \Omega_2$ . If  $F : \kappa \cap (\overline{\Omega}_2 \setminus \Omega_1) \rightarrow \kappa$  is a completely continuous operator such that, either

(i)  $\|Fx\| \leq \|x\|$ ,  $x \in \kappa \cap \partial\Omega_1$  and  $\|Fx\| \geq \|x\|$ ,  $x \in \kappa \cap \partial\Omega_2$ , or

(ii)  $\|Fx\| \geq \|x\|$ ,  $x \in \kappa \cap \partial\Omega_1$  and  $\|Fx\| \leq \|x\|$ ,  $x \in \kappa \cap \partial\Omega_2$  holds.

Then the operator  $F$  has a fixed point in  $\kappa \cap (\overline{\Omega}_2 \setminus \Omega_1)$ .

### 3. Existence of Positivity of the Solutions

This section presents the existence of positivity results of the problem stated in (1), (2).

For our construction, let us take  $X = \{x : x \in C[0, 1]\}$ , a Banach space with the standard norm  $\|x\| = \max_{t \in [0, 1]} |x(t)|$ . Consider a set  $\kappa$  as

$$\kappa = \{x \in X : x(t) \geq 0 \text{ for } t \in [0, 1] \text{ and } \min_{t \in [\xi, 1]} x(t) \geq \mathcal{M}\|x\|\},$$

where

$$\mathcal{M} = \min\{\xi^{m-1}, \xi^{n-1}\}. \quad (10)$$

Then, it is obvious that the set  $\kappa$  is a cone in  $X$ . To establish the results, we use the operator  $F : \kappa \rightarrow X$  by defining as

$$Fx(t) = \int_0^1 \overline{G}_m(t, s) \phi_q \left[ \int_0^1 G_n(s, r) g(r, x(r)) dr \right] ds. \quad (11)$$

Define the non-negative extended real numbers  $g_0$ ,  $g^0$ ,  $g_\infty$  and  $g^\infty$  by

$$g_0 = \lim_{x \rightarrow 0^+} \min_{t \in [0, 1]} \frac{g(t, x)}{\phi_p(x)}, \quad g^0 = \lim_{x \rightarrow 0^+} \max_{t \in [0, 1]} \frac{g(t, x)}{\phi_p(x)},$$

$$g_\infty = \lim_{x \rightarrow \infty} \min_{t \in [0, 1]} \frac{g(t, x)}{\phi_p(x)} \text{ and } g^\infty = \lim_{x \rightarrow \infty} \max_{t \in [0, 1]} \frac{g(t, x)}{\phi_p(x)},$$

and assume that the above are exist. The case  $g^0 = 0$  and  $g_\infty = \infty$  represents superlinear and the case  $g_0 = \infty$  and  $g^\infty = 0$  represents the sublinear.

We also consider the following conditions are fulfilled in this paper:

(F2)  $0 < \int_0^1 \overline{G}_m(t, s) ds < \infty$  and  $0 < \int_0^1 G_n(t, s) ds < \infty$ , and

(F3) the function  $g(t, x)$  is a non-decreasing for the second variable  $x$ .

**Lemma 3.1.** If  $F : \kappa \rightarrow X$  is given in (11) then  $F$  is a self map on the cone  $\kappa$ .

*Proof.* By using the Lemmas 2.3, 2.4 and the condition (F2),  $Fx(t) \geq 0$  for  $x \in \kappa$  and  $t \in [0, 1]$ . Then, by Lemma 2.3 and for  $x \in \kappa$ , we obtain

$$\begin{aligned} Fx(t) &= \int_0^1 \overline{G}_m(t, s) \phi_q \left( \int_0^1 G_n(s, r) g(r, x(r)) dr \right) ds \\ &\leq \int_0^1 \overline{G}_m(1, s) \phi_q \left( \int_0^1 G_n(s, r) g(r, x(r)) dr \right) ds \end{aligned}$$

so that

$$\|Fx\| \leq \int_0^1 \overline{G}_m(1, s) \phi_q \left( \int_0^1 G_n(s, r) g(r, x(r)) dr \right) ds. \quad (12)$$

Then, by Lemma 2.3 and (10), for  $x \in \kappa$  that

$$\begin{aligned} \min_{t \in [\xi, 1]} Fx(t) &= \min_{t \in [\xi, 1]} \left\{ \int_0^1 \overline{G}_m(t, s) \phi_q \left( \int_0^1 G_n(s, r) g(r, x(r)) dr \right) ds \right\} \\ &\geq \xi^{m-1} \int_0^1 \overline{G}_m(1, s) \phi_q \left( \int_0^1 G_n(s, r) g(r, x(r)) dr \right) ds \\ &\geq \mathcal{M} \int_0^1 \overline{G}_m(1, s) \phi_q \left( \int_0^1 G_n(s, r) g(r, x(r)) dr \right) ds \\ &\geq \mathcal{M} \|Fx\|. \end{aligned}$$

Hence, the operator  $F$  is a self map on a cone  $\kappa$ .  $\square$

Moreover by applying Arzela-Ascoli theorem,  $F$  is a completely continuous operator. The existence of positivity results of the  $p$ -Laplacian problem stated in (1), (2) for both the superlinear case as well as the sublinear case is now established.

**Theorem 3.2.** *Suppose the assumptions (F1), (F2) and (F3) are fulfilled. If the conditions  $g^0 = 0$  and  $g_\infty = \infty$  hold, then the nonlinear  $p$ -Laplacian problem stated in (1), (2) has at least one positive solution and it lies in the cone  $\kappa$ .*

*Proof.* From the definition of  $g^0 = 0$ , there exist  $\rho_1 > 0$  and  $\mathcal{H}_1 > 0$  such that

$$g(t, x) \leq \rho_1 \phi_p(x), \text{ for } 0 < x \leq \mathcal{H}_1,$$

where  $\rho_1$  satisfies

$$(\rho_1)^{q-1} \int_0^1 \overline{G}_m(1, s) \phi_q \left( \int_0^1 G_n(1, r) dr \right) ds \leq 1. \quad (13)$$

Let  $x \in \kappa$  and  $\|x\| = \mathcal{H}_1$ . Then, for  $t \in [0, 1]$  and by Lemmas 2.3, 2.4, we get

$$\begin{aligned} Fx(t) &= \int_0^1 \overline{G}_m(t, s) \phi_q \left( \int_0^1 G_n(s, r) g(r, x(r)) dr \right) ds \\ &\leq \int_0^1 \overline{G}_m(1, s) \phi_q \left( \int_0^1 G_n(1, r) \rho_1 \phi_p(x) dr \right) ds \\ &\leq (\rho_1)^{q-1} \int_0^1 \overline{G}_m(1, s) \phi_q \left( \int_0^1 G_n(1, r) dr \right) ds \|x\| \\ &\leq \|x\|. \end{aligned}$$

Hence,  $\|Fx\| \leq \|x\|$ . Now, if we are setting

$$\Omega_1 = \{x \in X : \|x\| < \mathcal{H}_1\}$$

then

$$\|Fx\| \leq \|x\|, \text{ for } x \in \kappa \cap \partial\Omega_1. \quad (14)$$

Next, since  $g_\infty = \infty$ , there exist  $\rho_2 > 0$  and  $\bar{\mathcal{H}}_2 > 0$  such that

$$g(t, x(t)) \geq \rho_2 \phi_p(x), \text{ for } x \geq \bar{\mathcal{H}}_2,$$

where  $\rho_2$  satisfies

$$(\rho_2)^{q-1} \mathcal{M}^2 \int_{s \in [\xi, 1]} \bar{G}_m(1, s) \phi_q \left( \mathcal{M} \int_{r \in [\xi, 1]} G_n(1, r) dr \right) ds \geq 1. \quad (15)$$

Let  $\mathcal{H}_2 = \max \left\{ 2\mathcal{H}_1, \frac{\bar{\mathcal{H}}_2}{\mathcal{M}} \right\}$ . Choose  $x \in \kappa$  and  $\|x\| = \mathcal{H}_2$ . Then

$$\min_{t \in [\xi, 1]} x(t) \geq \mathcal{M}\|x\| \geq \bar{\mathcal{H}}_2.$$

By the Lemmas 2.3, 2.4 and (10), and for  $t \in [0, 1]$ , we obtain

$$\begin{aligned} Fx(t) &= \int_0^1 \bar{G}_m(t, s) \phi_q \left( \int_0^1 G_n(s, r) g(r, x(r)) dr \right) ds \\ &\geq \min_{t \in [\xi, 1]} \left\{ \int_0^1 \bar{G}_m(t, s) \phi_q \left( \int_0^1 G_n(s, r) g(r, x(r)) dr \right) ds \right\} \\ &\geq \mathcal{M} \int_0^1 \bar{G}_m(1, s) \phi_q \left( \int_0^1 G_n(s, r) g(r, x(r)) dr \right) ds \\ &\geq \mathcal{M} \int_{s \in [\xi, 1]} \bar{G}_m(1, s) \phi_q \left( \mathcal{M} \int_{r \in [\xi, 1]} G_n(1, r) \rho_2 \phi_p(x) dr \right) ds \\ &\geq \mathcal{M} (\rho_2)^{q-1} \int_{s \in [\xi, 1]} \bar{G}_m(1, s) \phi_q \left( \mathcal{M} \int_{r \in [\xi, 1]} G_n(1, r) dr \right) \mathcal{M} \|x\| ds \\ &\geq (\rho_2)^{q-1} \mathcal{M}^2 \int_{s \in [\xi, 1]} \bar{G}_m(1, s) \phi_q \left( \mathcal{M} \int_{r \in [\xi, 1]} G_n(1, r) dr \right) \|x\| ds \\ &\geq \|x\|. \end{aligned}$$

Therefore,  $\|Fx\| \geq \|x\|$ . So, if we take

$$\Omega_2 = \{x \in X : \|x\| < \mathcal{H}_2\}$$

then

$$\|Fx\| \geq \|x\| \text{ for } x \in \kappa \cap \partial\Omega_2. \quad (16)$$

By Theorem 2.5 to (14) and (16), it follows that the operator  $F$  has a fixed point  $x \in \kappa \cap (\Omega_2 \setminus \bar{\Omega}_1)$  and that fixed point  $x$  is the positive solution of the  $p$ -Laplacian problem (1)-(2).  $\square$

**Theorem 3.3.** Suppose the assumptions (F1), (F2) and (F3) are fulfilled. If the conditions  $g_0 = \infty$  and  $g^\infty = 0$  hold, then the nonlinear  $p$ -Laplacian problem stated in (1), (2) has at least one positive solution and it lies in the cone  $\kappa$ .

*Proof.* From the definition of  $g_0 = \infty$ , there exist  $\bar{\rho}_1 > 0$  and  $\mathcal{R}_1 > 0$  such that

$$g(t, x) \geq \bar{\rho}_1 \phi_p(x), \text{ for } 0 < x \leq \mathcal{R}_1,$$

where  $\bar{\rho}_1 \geq \rho_2$  and  $\rho_2$  is given in (15).

Let  $x \in \kappa$  and  $\|x\| = \mathcal{R}_1$ . Then, by Lemmas 2.3, 2.4 and (10), and for  $t \in [0, 1]$ , we obtain

$$\begin{aligned} Fx(t) &= \int_0^1 \overline{G}_m(t, s) \phi_q \left( \int_0^1 G_n(s, r) g(r, x(r)) dr \right) ds \\ &\geq \min_{t \in [\xi, 1]} \left\{ \int_0^1 \overline{G}_m(t, s) \phi_q \left( \int_0^1 G_n(s, r) g(r, x(r)) dr \right) ds \right\} \\ &\geq \mathcal{M} \int_{s \in [\xi, 1]} \overline{G}_m(1, s) \phi_q \left( \int_{r \in [\xi, 1]} G_n(s, r) \bar{\rho}_1 \phi_p(x) dr \right) ds \\ &\geq \mathcal{M} \int_{s \in [\xi, 1]} \overline{G}_m(1, s) \phi_q \left( \mathcal{M} \int_{r \in [\xi, 1]} G_n(1, r) \bar{\rho}_1 \phi_p(x) dr \right) ds \\ &\geq \mathcal{M} (\bar{\rho}_1)^{q-1} \int_{s \in [\xi, 1]} \overline{G}_m(1, s) \phi_q \left( \mathcal{M} \int_{r \in [\xi, 1]} G_n(1, r) dr \right) \mathcal{M} \|x\| ds \\ &= (\bar{\rho}_1)^{q-1} \mathcal{M}^2 \int_{s \in [\xi, 1]} \overline{G}_m(1, s) \phi_q \left( \mathcal{M} \int_{r \in [\xi, 1]} G_n(1, r) dr \right) \|x\| ds \\ &\geq \|x\|. \end{aligned}$$

Therefore,  $\|Fx\| \geq \|x\|$ . Now, if we take

$$\Omega_3 = \{x \in X : \|x\| < \mathcal{R}_1\}$$

then

$$\|Fx\| \geq \|x\|, \text{ for } x \in \kappa \cap \partial\Omega_3. \quad (17)$$

Next, since  $g^\infty = 0$ , there exist  $\bar{\rho}_2 > 0$  and  $\bar{\mathcal{R}}_2 > 0$  such that

$$g(t, x(t)) \leq \bar{\rho}_2 \phi_p(x), \text{ for } x \geq \bar{\mathcal{R}}_2,$$

where  $\bar{\rho}_2 \leq \rho_1$  and  $\rho_1$  is given in (13).

Set

$$g^*(t, x) = \sup_{0 \leq s \leq x} g(t, s).$$

Then, it is evident that the real-valued function  $g^*$  is a non-decreasing,  $g \leq g^*$  and

$$\lim_{x \rightarrow \infty} \frac{g^*(t, x)}{x} = 0.$$

It follows that there exists  $\mathcal{R}_2 > \max\{2\mathcal{R}_1, \bar{\mathcal{R}}_2\}$  such that

$$g^*(t, x) \leq g^*(t, \mathcal{R}_2), \text{ for } 0 < x \leq \mathcal{R}_2.$$

Choose  $x \in \kappa$  with  $\|x\| = \mathcal{R}_2$ . Then, we get

$$\begin{aligned} Fx(t) &= \int_0^1 \overline{G}_m(t, s) \phi_q \left( \int_0^1 G_n(s, r) g(r, x(r)) dr \right) ds \\ &\leq \int_0^1 \overline{G}_m(1, s) \phi_q \left( \int_0^1 G_n(s, r) g(r, \mathcal{J}_2) dr \right) ds \\ &\leq \int_0^1 \overline{G}_m(1, s) \phi_q \left( \int_0^1 G_n(s, r) \bar{\rho}_2 \phi_p(\mathcal{J}_2) dr \right) ds \\ &\leq (\bar{\rho}_2)^{q-1} \int_0^1 \overline{G}_m(1, s) \phi_q \left( \int_0^1 G_n(1, r) dr \right) ds \mathcal{R}_2 \\ &\leq \mathcal{R}_2 = \|x\|. \end{aligned}$$

Hence,  $\|Fx\| \leq \|x\|$ . Now, if we take

$$\Omega_4 = \{x \in X : \|x\| < \mathcal{R}_2\}$$

then

$$\|Fx\| \leq \|x\|, \text{ for } x \in \kappa \cap \partial\Omega_4. \quad (18)$$

Applying by Theorem 2.5 to (17) and (18), we have that the operator  $F$  has a fixed point  $x \in \kappa \cap (\Omega_4 \setminus \bar{\Omega}_3)$  and that fixed point  $x$  is the positive solution of the  $p$ -Laplacian problem (1)-(2).  $\square$

We consider the example to validate the established results.

**Example 3.4.** Let us take  $m = 4$ ,  $n = 3$  and  $\xi = \frac{1}{3}$ . Now, consider a nonlinear  $p$ -Laplacian problem

$$[\phi_p(x^{(4)}(t))]^{(3)} = g(t, x(t)), \quad t \in [0, 1], \quad (19)$$

satisfying

$$\left. \begin{aligned} x(0) = 0, \quad x^{(1)}(0) = 0, \quad x^{(2)}(0) = 0, \quad x^{(2)}(1) - 2x^{(2)}\left(\frac{1}{3}\right) &= 0, \\ [\phi_p(x^{(4)}(t))]_{\text{at } t=0} = 0, \quad [\phi_p(x^{(4)}(t))]_{\text{at } t=0}' &= 0, \\ [\phi_p(x^{(4)}(t))]_{\text{at } t=1}' - 2[\phi_p(x^{(4)}(t))]_{\text{at } t=\xi=\frac{1}{3}}' &= 0. \end{aligned} \right\} \quad (20)$$

Let us take  $p = 2$  for simplicity. By algebraic computations, we get

$$\mathcal{M} = 0.1111.$$

- (a) If we take  $g(t, x(t)) = x^2(1 - e^{-2t})$ , then  $g^0 = 0$  and  $g_\infty = \infty$ . So, all the claims in the Theorem 3.2 are fulfilled. Therefore, the boundary value problem (19)-(20) has at least one positive solution.
- (b) If we take  $g(t, x(t)) = \frac{\sqrt{(t^2+1)}}{t^2x^2}$ , then  $g_0 = \infty$  and  $g^\infty = 0$ . So, all the claims in the Theorem 3.3 are fulfilled. Therefore, the boundary value problem (19)-(20) has at least one positive solution.

#### 4. Conclusion

In this paper, we established the existence of positive solutions to the higher order three-point boundary value problem involving  $p$ -Laplacian operator by an application of the Guo–Krasnosel'skii fixed point theorem for operators on a cone in a Banach space.

**Acknowledgements:** The authors express their gratitude to the referees for their insightful recommendations and remarks.

#### References

- [1] H. Afshari, S. Kalantari, E. Karapinar, Solution of fractional differential equations via coupled fixed point, Electron. J. Differ. Equ. 2015 (2015) 1-12.
- [2] H. Afshari, E. Karapinar, A discussion on the existence of positive solutions of the boundary value problems via  $\psi$ -Hilfer fractional derivative on  $b$ -metric spaces, Adv. Differ. Equ. 2020 (2020) 1-11.
- [3] R.P. Agarwal, H. Lü, D. O'Regan, Eigenvalues and the one-dimensional  $p$ -Laplacian, J. Math. Anal. Appl. 266 (2002) 383-400.
- [4] R.I. Avery, J. Henderson, Existence of three positive pseudo-symmetric solutions for a one-dimensional  $p$ -Laplacian, J. Math. Anal. Appl. 277 (2003) 395-404.
- [5] A.C. Cavalcione, Existence results for Navier problems with degenerated  $(p, q)$ -Laplacian and  $(p, q)$ -Biharmonic operators, Results Nonlinear Anal. 1 (2018) 74-87.
- [6] K. Deimling, Nonlinear Functional Analysis, Springer-Verlag, New York, 1985.

- [7] L. Diening, P. Lindqvist, B. Kawohl, Mini-Workshop: The  $p$ -Laplacian Operator and Applications, Oberwolfach Reports, Report No. 08/2013, 433–482.
- [8] Y. Ding, J. Xu, X. Zhang, Positive solutions for a  $2n^{\text{th}}$  order  $p$ -Laplacian boundary value problem involving all derivatives, Electron. J. Differ. Equ. 2013 (2013) 1–14.
- [9] G. Dwivedi, Generalised Picone's identity and some qualitative properties of  $p$ -sub-Laplacian on Heisenberg group, Adv. Theory Nonlinear Anal. Appl. 5 (2) (2021) 232–239.
- [10] H. Feng, H. Pang, W. Ge, Multiplicity of symmetric positive solutions for a multi-point boundary value problem with a one-dimensional  $p$ -Laplacian, Nonlinear Anal. 69 (2008) 3050–3059.
- [11] J.R. Graef, B. Yang, Multiple positive solutions to a three-point third order boundary value problem, Discrete Contin. Dyn. Syst. 2005 (2005) (Special) 337–344.
- [12] Y. Guo, W. Ge, Twin positive symmetric solutions for Lidstone boundary value problems, Taiwan. J. Math. 8 (2) (2004) 271–283.
- [13] Y. Guo, Y. Ji, X. Liu, Multiple positive solutions for some multi-point boundary value problems with  $p$ -Laplacian, J. Comput. Appl. Math. 216 (2008) 144–156.
- [14] D. Guo, V. Lakshmikantham, Nonlinear Problems in Abstract Cones, Academic Press, San Diego, CA, 1988.
- [15] E. Karapinar, H.D. Binh, N.H. Luc, N.H. Can, On continuity of the fractional derivative of the time-fractional semilinear pseudo-parabolic systems, Adv. Differ. Equ. 2021 (2021) 1–24.
- [16] E. Karapinar, A. Fulga, M. Rashid, L. Shahid, H. Aydi, Large contractions on quasi-metrics spaces with an application to nonlinear fractional differential equations, Mathematics 7 (5) (2019) 1–11.
- [17] M. Krasnoselskii, Positive Solutions of Operator Equations, Noordhoff, Groningen, 1964.
- [18] J.E. Lazreg, S. Abbas, M. Benchohra, E. Karapinar, Impulsive Caputo-Fabrizio fractional differential equations in  $b$ -metric spaces, Open Mathematics 19 (1) (2021) 363–372.
- [19] C. Li, W. Ge, Existence of positive solutions for  $p$ -Laplacian singular boundary value problems, Indian J. Pure Appl. Math. 34 (2003) 187–203.
- [20] J. Li, J. Shen, Existence of three positive solutions for boundary value problems with  $p$ -Laplacian, J. Math. Anal. Appl. 311 (2005) 457–465.
- [21] W.C. Lian, F.H. Wong, Existence of positive solutions for higher order generalized  $p$ -Laplacian BVPs, Appl. Math. Lett. 13 (2000) 35–43.
- [22] J. Liang, Z.W. Lv, Solutions to a three-point boundary value problem, Adv. Differ. Equ. 2011 (2011) 1–20.
- [23] X. Lin, Z. Fu, Positive solutions for a class of third order three-point boundary value problem, Discrete Dyn. Nat. Soc. 2012 (2012) 1–12.
- [24] X. Lin, Z. Zhao, Iterative technique for a third order differential equation with three-point nonlinear boundary value conditions, Electron. J. Qual. Theory Differ. Equ. 2016 (2016) 1–10.
- [25] Z. Liu, H. Chen, C. Liu, Positive solutions for singular third order non-homogeneous boundary value problems, J. Appl. Math. Comput. 38 (2012) 161–172.
- [26] Y. Liu, W. Ge, Multiple positive solutions to a three-point boundary value problem with  $p$ -Laplacian, J. Math. Anal. Appl. 277 (2003) 293–302.
- [27] D. Liu, Z. Ouyang, Solvability of third order three-point boundary value problems, Abstr. Appl. Anal. 2014 (2014) 1–7.
- [28] A. Ourraoui, Existence and uniqueness of solutions for Steklov problem with variable exponent, Adv. Theory Nonlinear Anal. Appl. 5 (1) (2021) 158–166.
- [29] A.P. Palamides, G. Smyrdis, Positive solutions to a singular third order three-point boundary value problem with an indefinitely signed Green's function, Nonlinear Anal. TMA, 68 (7) (2008) 2104–2118.
- [30] K.R. Prasad, N. Sreedhar, L.T. Wesen, Existence of positive solutions for higher order  $p$ -Laplacian boundary value problems, Mediterr. J. Math. 15 (2018) 1–12.
- [31] R.R. Sankar, N. Sreedhar, K.R. Prasad, Existence of positive solutions for  $3n^{\text{th}}$  order boundary value problems involving  $p$ -Laplacian, Creat. Math. Inform. 31 (1) (2022) 101–108.
- [32] G. Shi, J. Zhang, Positive solutions for higher order singular  $p$ -Laplacian boundary value problems, Proc. Indian Acad. Sci. Math. Sci. 118 (2008) 295–305.
- [33] Y. Sun, Positive solutions for third order three-point non-homogeneous boundary value problems, Appl. Math. Lett. 22 (2009) 45–51.
- [34] Y. Sun, Q. Sun, X. Zhang, Existence and non existence of positive solutions for a higher order three-point boundary value problem, Abstr. Appl. Anal. 2014 (2014) 1–7.
- [35] C.X. Wang, H.R. Sun, Positive solutions for a class of singular third order three-point non-homogeneous boundary value problem, Dynam. Syst. Appl. 19 (2010) 225–234.
- [36] Z.L. Wei, Existence of positive solutions for  $n^{\text{th}}$  order  $p$ -Laplacian singular sublinear boundary value problems, Appl. Math. Lett. 36 (2014) 25–30.
- [37] Z.L. Wei, Existence of positive solutions for  $n^{\text{th}}$  order  $p$ -Laplacian singular super-linear boundary value problems, Appl. Math. Lett. 50 (2015) 133–140.
- [38] J. Xu, Z. Wei, Y. Ding, Positive solutions for a  $2n^{\text{th}}$  order  $p$ -Laplacian boundary value problem involving all even derivatives, Topol. Method Nonlinear Anal. 39 (2012) 23–36.
- [39] L. Zhao, W. Wang, C. Zhai, Existence and uniqueness of monotone positive solutions for a third order three-point boundary value problem, Differ. Equ. Appl. 10 (3) (2018) 251–260.